arXiv:hep-ph/9609447v4 18 Nov 1996

hep-ph/9609447
Freiburg–THEP 96/19
September 1996

Analytic evaluation of two–loop renormalization constants of enhanced electroweak strength in the Higgs sector of the Standard Model

V. Borodulin

Institute for High Energy Physics
Protvino, Moscow Region 142284, Russia

G. Jikia

Albert–Ludwigs–Universität Freiburg, Fakultät für Physik
Hermann–Herder Str.3, D-79104 Freiburg, Germany

Abstract

We calculate renormalization constants $Z_{H,w}$, the Higgs and W–boson mass and tadpole counterterms in the on–mass-shell renormalization scheme to two loops in the heavy–Higgs–boson limit $m_H \gg M_W$. Explicit analytic formulae are presented for the two–loop integrals with masses, which are not known in the literature. As an application, the analytic expression for the two–loop leading correction to the fermionic Higgs boson width is obtained.

1. Introduction

After the observation of the top quark signal at the Tevatron, the mechanism of the spontaneous electroweak symmetry breaking remains the last untested property of the Standard Model ($\mathcal{SM}$). Although the recent global fits to the precision electroweak data from LEP, SLC and Tevatron seem to indicate a preference to a light Higgs boson $m_H = 149^{+148}_{-82}$ GeV, $m_H < 450$ GeV ($95\% C.L.$) [1], it is definitely premature to exclude the heavy Higgs scenario. The reason is that a restrictive upper bound for m_H is dominated by the result on A_{LR}, which differs significantly from the $\mathcal{SM}$ predictions [2]. Without A_{LR} upper bound on M_H becomes larger than 600 GeV [2], which is not far in the logarithmic scale from the value of the order of 1 TeV, where perturbation theory breaks down. In order to estimate the region of applicability of the perturbation theory, the leading two–loop $\mathcal{O}(g^4 m_H^4/M_W^4)$ electroweak corrections were under intense study recently. In particular, the high energy weak–boson scattering [3], corrections to the heavy Higgs line shape [4], corrections to the partial widths of the Higgs boson decay to pairs of fermions [5, 6] and intermediate vector bosons [7, 8] at two–loops have been calculated. All these calculations resort at least partly to numerical methods. Even for the two–loop renormalization constants in the Higgs scalar sector of the $\mathcal{SM}$ [4, 9] complete analytical expressions are not known. In this paper we present our analytic results for these two–loop renormalization constants, evaluated in the on–mass–shell renormalization scheme in terms of transcendental functions $\zeta(3)$ and the maximal value of the Clausen function $\mathrm{Cl}(\pi/3)$.

2. Lagrangian and renormalization

The part of the $\mathcal{SM}$ Lagrangian describing the Higgs scalar sector of the $\mathcal{SM}$ in terms of the bare quantities is given by:

$$\begin{aligned}\mathcal{L} = &\frac{1}{2}\partial_\mu H_0 \partial^\mu H_0 + \frac{1}{2}\partial_\mu z_0 \partial^\mu z_0 + \partial_\mu w_0^+ \partial^\mu w_0^- \\ &- \frac{m_{H0}^2}{2v_0^2}\left(w_0^+ w_0^- + \frac{1}{2}z_0^2 + \frac{1}{2}H_0^2 + v_0 H_0 + \frac{1}{2}\delta v^2\right)^2 .\end{aligned} \tag{2.1}$$

Here the tadpole counterterm δv^2 is chosen in such a way, that a Higgs field vacuum expectation value is equal to v_0

$$v_0 = \frac{2\, M_{W0}}{g} \tag{2.2}$$

to all orders.

Renormalized fields are given by

$$H_0 = \sqrt{Z_H}\, H, \quad z_0 = \sqrt{Z_z}\, z, \quad w_0 = \sqrt{Z_w}\, w. \tag{2.3}$$

At two–loop approximation the wave function renormalization constants, tadpole and mass counterterms take the form

$$\begin{aligned}\sqrt{Z} &= 1 + \frac{g^2}{16\pi^2}\,\delta Z^{(1)} + \frac{g^4}{(16\pi^2)^2}\,\delta Z^{(2)}; \\ \delta v^2 &= \frac{1}{16\pi^2}\,\delta v^{2\,(1)} + \frac{g^2}{(16\pi^2)^2}\,\delta v^{2\,(2)};\end{aligned} \tag{2.4}$$

$$
\begin{aligned}
M_{W0} &= M_W + \frac{g^2}{16\pi^2}\,\delta M_W^{(1)} + \frac{g^4}{(16\pi^2)^2}\,\delta M_W^{(2)};\\
m_{H0}^2 &= m_H^2 + \frac{g^2}{16\pi^2}\,\delta m_H^2{}^{(1)} + \frac{g^4}{(16\pi^2)^2}\,\delta m_H^2{}^{(2)}.
\end{aligned}
$$

Since weak coupling constant g is not renormalized at the leading order in m_H^2, the W–boson mass counterterm is related to the Nambu–Goldstone wave function renormalization constant by the Ward identity $M_{W0} = Z_w M_W$.

In the on–mass–shell renormalization scheme all the counterterms are fixed uniquely by the requirement that the pole position of the Higgs and W–boson propagators coincide with their physical masses and the residue of the Higgs boson pole is normalized to unity.

The one–loop counterterms equivalent to those used in [4, 9] are given by

$$
\begin{aligned}
\delta v^2{}^{(1)} &= m_H^2 \xi_H^\epsilon \left\{ -\frac{6}{\epsilon} + 3 - \epsilon\Big(\frac{3}{2} + \frac{\pi^2}{8}\Big)\right\};\\
\frac{\delta m_H^2{}^{(1)}}{m_H^2} &= \frac{m_H^2 \xi_H^\epsilon}{M_W^2}\left\{ -\frac{3}{\epsilon} + 3 - \frac{3\,\pi\,\sqrt{3}}{8} + \epsilon\Big(-3 + \frac{\pi^2}{16} + \frac{3\pi\sqrt{3}}{8} + \frac{3\sqrt{3}C}{4} - \frac{3\pi\sqrt{3}\log 3}{16}\Big)\right\};\\
\delta Z_H^{(1)} &= \frac{m_H^2 \xi_H^\epsilon}{M_W^2}\left\{ \frac{3}{4} - \frac{\pi\,\sqrt{3}}{8} + \epsilon\Big(-\frac{3}{4} + \frac{3\pi\sqrt{3}}{32} + \frac{\sqrt{3}C}{4} - \frac{\pi\sqrt{3}\log 3}{16}\Big)\right\};\\
\frac{\delta M_W^{(1)}}{M_W} &= \delta Z_w^{(1)} = -\frac{m_H^2 \xi_H^\epsilon}{16\,M_W^2}\left\{1 - \frac{3}{4}\epsilon\right\}.
\end{aligned}
\tag{2.5}
$$

Here the dimension of space–time is taken to be $d = 4 + \epsilon$ and

$$
\xi_H^\epsilon = e^{\gamma\,\epsilon/2}\left(\frac{m_H}{2\,\pi}\right)^{\epsilon}. \tag{2.6}
$$

In contrast to papers [4, 9] we prefer not to keep the one–loop counterterms of $\mathcal{O}(\epsilon)$ order, *i.e.* unlike in the conventional on–mass–shell scheme used in [4, 9], we require that the one-loop normalization conditions are fulfilled only in the limit $\epsilon \to 0$, where the counterterms of the order $\mathcal{O}(\epsilon)$ do not contribute. Such a modified on–mass-shell scheme is equally consistent as the conventional one or the standard scheme of minimal dimensional renormalization, which assumes only the subtraction of pole terms at $\epsilon = 0$ [10]. (Moreover, in general one cannot subtract all the nonsingular $\mathcal{O}(\epsilon)$ terms in the Laurent expansion in ϵ, as they are not polynomial in external momenta.) The $\mathcal{O}(\epsilon)$ one–loop counterterms considered in [4, 9] do can really combine with the $1/\epsilon$ terms at two–loop order to give finite contributions, but these contributions are completely canceled by the additional finite parts of the two–loop counterterms, fixed through the renormalization conditions in the on–mass–shell renormalization scheme. The reason is that after the inclusion of the one–loop counterterms all the subdivergences are canceled and only the overall divergence remains, which is to be canceled by the two–loop counterterms. The account of finite contributions coming from the combination of $\mathcal{O}(\epsilon)$ one–loop counterterms with $1/\epsilon$ overall divergence just redefines the finite parts of the two–loop counterterms. An obvious advantage of this modified on–mass–shell scheme is that the lower loop counterterms once calculated could be directly used in higher loop calculations, while in the conventional on–mass–shell scheme for l-loop calculation one needs to recalculate the one-loop counterterms to include all the terms up to $\mathcal{O}(\epsilon^{l-1})$, two-loop counterterms to include $\mathcal{O}(\epsilon^{l-2})$ terms and so on.

3. Analytic integration

The calculation of the Higgs and W–boson (or Nambu–Goldstone w, z bosons) two–loop self energies, needed to evaluate the renormalization constants (2.4), reduces to the evaluation of the two–loop massive scalar integrals

$$J(k^2; n_1\, m_1^2, n_2\, m_2^2, n_3\, m_3^2, n_4\, m_4^2, n_5\, m_5^2) = -\frac{1}{\pi^4}\int D^{(d)}P\, D^{(d)}Q\, \left(P^2 - m_1^2\right)^{-n_1} \tag{3.1}$$
$$\times\left((P+k)^2 - m_2^2\right)^{-n_2}\left((Q+k)^2 - m_3^2\right)^{-n_3}\left(Q^2 - m_4^2\right)^{-n_4}\left((P-Q)^2 - m_5^2\right)^{-n_5}$$

and their derivatives J' at $k^2 = m_H^2$ or at $k^2 = 0$.

The most difficult is a calculation of the all–massive scalar master integral corresponding to the topology shown in the Fig. 1.



Fig. 1. The two loop all–massive master graph. Solid line represents Higgs bosons.

This integral has a discontinuity that is an elliptic integral, resulting from integration over the phase space of three massive particles, and is not expressible in terms of polylogarithms. However one can show [11] that on–shell $k^2 = m_H^2$ or at the threshold $k^2 = 9m_H^2$ this is not the case. We use the dispersive method [12, 13] to evaluate this finite integral on the mass shell:

$$m_H^2\, J(k^2; m_H^2, m_H^2, m_H^2, m_H^2, m_H^2) = \sigma_a(k^2/m_H^2) + \sigma_b(k^2/m_H^2), \tag{3.2}$$

where $\sigma_{a,b}$ correspond to the dispersive integrals calculated, respectively, from the two– and three–particle discontinuities, which are itself reduced to one–dimensional integrals. The $\tanh^{-1}$ functions entering $\sigma_{a,b}$ can be removed integrating by parts either in the dispersive integral [12], or in the discontinuity integral [13]. By interchanging the order of integrations the latter representation gives the three–particle cut contribution σ_b as a single integral of logarithmic functions [13]. After some rather heavy analysis we obtain at $k^2 = m_H^2$:

$$\begin{aligned}
\sigma_a(1) &= \int_0^1 dy\, \frac{8}{y^4 - y^2 + 1}\log\left(\frac{(y^2+1)^2}{y^4+y^2+1}\right)\left[\frac{(y^4-1)\log(y)}{y} - \frac{\pi\, y}{\sqrt{3}}\right] \\
&= \frac{17}{18}\zeta(3) - \frac{10}{9}\pi\, C + \pi^2\log 2 - \frac{4}{9}\pi^2\log 3, &(3.3)\\
\sigma_b(1) &= \int_0^1 dy\, 2\log\left(\frac{y^2+y+1}{y}\right) \\
&\left[\frac{\log(y+1)}{y} + \frac{\frac{\pi}{\sqrt{3}}(y^2 - 3\,y + 1) - 2\,(y^2-1)\log(y)}{y^4+y^2+1} - \frac{\log(y)}{y+1}\right] \\
&= \frac{1}{18}\zeta(3) + \frac{4}{9}\pi\, C - \pi^2\log 2 + \frac{4}{9}\pi^2\log 3. &(3.4)
\end{aligned}$$

Here $C = \mathrm{Cl}(\pi/3) = \mathrm{Im}\ \mathrm{li}_2\left(\exp\left(i\pi/3\right)\right) = 1.01494\,16064\,09653\,62502\ldots$

As a result we find

$$\begin{aligned} m_H^2\, J(m_H^2; m_H^2, m_H^2, m_H^2, m_H^2, m_H^2) &= \zeta(3) - \frac{2}{3}\pi C \\ &= -0.92363\,18265\,19866\,53695\ldots \end{aligned} \tag{3.5}$$

The numerical value is in agreement with the one, calculated using the momentum expansion [14], and with the numerical values given in [9, 15].

Given the value (3.5), the simplest way to calculate the derivative of the integral (3.2) is to use Kotikov's method of differential equations [16, 9]

$$\begin{aligned} m_H^4\, J'(m_H^2; m_H^2, m_H^2, m_H^2, m_H^2, m_H^2) &= \frac{2}{3}\pi C - \zeta(3) - \frac{\pi^2}{9} \\ &= -0.17299\,08847\,12284\,42069\ldots \end{aligned} \tag{3.6}$$

All the other two–loop self energy scalar integrals contain "light" particles (Nambu–Goldstone or W, Z–bosons) and some of them are IR divergent in the limit $M_{W,Z} \to 0$. In principle, one can calculate these IR divergent integrals in Landau gauge, where masses of Nambu–Goldstone bosons are equal to zero and IR divergences are represented as (double) poles at $\epsilon = 0$ [9]. However, in order to have an additional check of the cancellation in the final answer of all the IR divergent $\log(M_{W,Z}^2)$–terms, we work in 't Hooft–Feynman gauge. For the infra–red finite integrals the correct answer in the leading order in m_H^2 is obtained just by setting $M_{W,Z} = 0$. We agree with the results for these integrals given in [9, 11]. The two–loop IR divergent integrals correspond to the topologies shown in the Fig. 2, which contain "massless" propagators squared.



Fig. 2. The two–loop IR divergent graphs. Dashed line represents "light" particles.

The relevant technique to handle these diagrams follows from the so–called asymptotic operation method [17]. According to the recipe of As–operation, the formal Taylor expansion in small mass M_W entering propagator (or its powers) should be accomplished by adding the terms, containing the δ–function or its derivatives. The additional terms counterbalance the infra–red singularities, arising in the formal expansion of propagator. In our case we have

$$\begin{aligned} \frac{1}{(P^2 - M_W^2)^2} &= \frac{1}{(P^2)^2} + 2\frac{M_W^2}{(P^2)^3} + \ldots \\ &+ C_1(M_W)\delta^{(d)}(P) + C_2(M_W)\partial^2\delta^{(d)}(P) + \ldots \end{aligned} \tag{3.7}$$

Here the first coefficient functions $C_i(M_W)$ read

$$\begin{aligned} C_1(M_W) &= \int D^{(d)}P \frac{1}{(P^2 - M_W^2)^2} \sim \mathcal{O}(M_W^0), \\ C_2(M_W) &= \frac{1}{2d}\int D^{(d)}P \frac{P^2}{(P^2 - M_W^2)^2} \sim \mathcal{O}(M_W^2). \end{aligned} \tag{3.8}$$

This equality is to be understood in the following sense [17]. One should integrate both parts of the equation multiplied by a test function and then take the limit $d \to 4$. If one keeps in the expansion all terms up to order M_W^{2n}, the resulting expression will represent a correct expansion of the initial integral to order $o(M_W^{2n})$. To obtain the leading contribution to the diagrams Fig. 2, it suffices just to take the first term of the Taylor expansion and, correspondingly, the first "counterterm" $C_1 \delta^{(d)}(P)$.

Finally, the combination of Mellin–Barnes representation and Kotikov's method gives the following answer, corresponding to the first graph in Fig. 2 neglecting the terms of order $\mathcal{O}(M_W^2/m_H^2)$:

$$
\begin{aligned}
m_H^2\, J(m_H^2; 2\,M_W^2, M_W^2, 0, M_W^2, m_H^2) &= \xi_H^{2\epsilon}\Bigg(\frac{2\,i\,\pi}{\epsilon} - \frac{i\,\pi}{2} + \frac{2}{\epsilon}\,\log(\frac{M_W^2}{m_H^2}) \\
&\quad -\frac{1}{2} + \frac{5\,\pi^2}{6} - \log(\frac{M_W^2}{m_H^2}) + \frac{1}{2}\log^2(\frac{M_W^2}{m_H^2})\Bigg); \qquad (3.9)\\
m_H^4\, J'(m_H^2; 2\,M_W^2, M_W^2, 0, M_W^2, m_H^2) &= \xi_H^{2\epsilon}\Bigg(-\frac{2\,i\,\pi}{\epsilon} + 2\,i\,\pi - \frac{2}{\epsilon}\left(1 \,+\, \log(\frac{M_W^2}{m_H^2})\right) \qquad (3.10)\\
&\quad +2 - \frac{5\,\pi^2}{6} + \log(\frac{M_W^2}{m_H^2}) - \frac{1}{2}\log^2(\frac{M_W^2}{m_H^2})\Bigg).
\end{aligned}
$$

The integral diverges as $1/\epsilon$, while if we would set $M_W = 0$ from the very beginning, it would diverge as $1/\epsilon^2$. The integral corresponding to the second graph in the Fig. 2 up to $\mathcal{O}(M_W^2/m_H^2)$ is

$$
\begin{aligned}
J(m_H^2; 2\,M_W^2, 0, 0, M_W^2, m_H^2) &= \xi_H^{2\epsilon}\Bigg(\frac{2}{\epsilon^2} + \frac{1}{\epsilon}\left(2\,\log(\frac{M_W^2}{m_H^2}) - 1\right) \qquad (3.11)\\
&\quad +\frac{1}{2} - \frac{\pi^2}{12} - \log(\frac{M_W^2}{m_H^2}) + \frac{1}{2}\log^2(\frac{M_W^2}{m_H^2})\Bigg)
\end{aligned}
$$

The two–loop vacuum integrals needed to evaluate the tadpole counterterm $\delta v^{2\,(2)}$ have been calculated in [18].

4. Results

The analytic results for the two–loop renormalization constants are:

$$
\begin{aligned}
\delta v^{2\,(2)} &= \frac{m_H^4 \xi_H^{2\epsilon}}{16\,M_W^2}\left(\frac{72}{\epsilon^2} + \frac{36\,\pi\,\sqrt{3} - 84}{\epsilon} - 162 - 3\,\pi^2 + 60\,\sqrt{3}C\right); \qquad (4.1)\\
\frac{\delta m_H^{2\;(2)}}{m_H^2} &= \frac{m_H^4 \xi_H^{2\epsilon}}{64\,M_W^4}\Bigg(\frac{576}{\epsilon^2} + \frac{144\,\pi\,\sqrt{3} - 1014}{\epsilon} \\
&\quad +\frac{99}{2} - 252\,\zeta(3) + 87\,\pi^2 - 219\,\pi\,\sqrt{3} \\
&\quad +156\,\pi\,C + 204\,\sqrt{3}C\Bigg); \qquad (4.2)\\
\delta Z_H^{(2)} &= \frac{m_H^4 \xi_H^{2\epsilon}}{64\,M_W^4}\Bigg(\frac{3}{\epsilon} - \frac{75}{4} - 126\,\zeta(3)
\end{aligned}
$$

$$+\frac{25\,\pi^2}{2} - 76\,\pi\,\sqrt{3} + 78\,\pi\,C + 108\,\sqrt{3}C\Bigg); \tag{4.3}$$

$$\frac{\delta M_W^{(2)}}{M_W} \;=\; \delta Z_w^{(2)} = \frac{m_H^4 \xi_H^{2\epsilon}}{64\,M_W^4}\left(\frac{3}{\epsilon} - \frac{3}{8} - \frac{\pi^2}{6} + \frac{3\,\pi\,\sqrt{3}}{2} - 6\,\sqrt{3}C\right). \tag{4.4}$$

For comparison we have also calculated these counterterms following the renormalization scheme [4, 5] and keeping the $\mathcal{O}(\epsilon)$ terms and found complete agreement with their (partly numerical) results. In this scheme the renormalization constants (4.1), (4.2) look a bit more complicated due to the presence of the additional $\pi\sqrt{3}\log 3$ terms

$$\begin{aligned}
\delta v^{2\,(2)} &= \frac{m_H^4 \xi_H^{2\epsilon}}{16\,M_W^2}\Bigg(\frac{72}{\epsilon^2} + \frac{36\,\pi\,\sqrt{3} - 84}{\epsilon} + 90 - 12\,\pi^2 - 12\,\sqrt{3}C \\
&\quad -36\,\pi\,\sqrt{3} + 18\,\pi\,\sqrt{3}\,\log 3\Bigg);
\end{aligned} \tag{4.5}$$

$$\begin{aligned}
\frac{\delta m_H^{2\;(2)}}{m_H^2} &= \frac{m_H^4 \xi_H^{2\epsilon}}{64\,M_W^4}\Bigg(\frac{576}{\epsilon^2} + \frac{144\,\pi\,\sqrt{3} - 1014}{\epsilon} \\
&\quad +\frac{2439}{2} - 252\,\zeta(3) + 63\,\pi^2 - 363\,\pi\,\sqrt{3} \\
&\quad +156\,\pi\,C - 84\,\sqrt{3}C + 72\,\pi\sqrt{3}\,\log 3\Bigg).
\end{aligned} \tag{4.6}$$

The wave function renormalization constants $Z_{H,w}$ are identical in these two schemes.

As an example of the physical quantity, for which all the schemes should give the same result, we consider the two–loop heavy Higgs correction to the fermionic Higgs width [5, 6]. The correction is given by the ratio

$$\begin{aligned}
\frac{Z_H}{M_{W0}^2/M_W^2} &= 1 + 2\frac{g^2}{16\,\pi^2}\left(\delta Z_H^{(1)} - \frac{\delta M_W^{(1)}}{M_W}\right) \\
&\quad +\frac{g^4}{(16\,\pi^2)^2}\Bigg[2\,\frac{\delta M_W^{(1)}}{M_W}\left(\frac{\delta M_W^{(1)}}{M_W} - \delta Z_H^{(1)}\right) + \left(\delta Z_H^{(1)} - \frac{\delta M_W^{(1)}}{M_W}\right)^2 \\
&\quad +2\,\delta Z_H^{(2)} - 2\,\frac{\delta M_W^{(2)}}{M_W}\Bigg].
\end{aligned} \tag{4.7}$$

Substituting (2.5), (4.3)–(4.4) we find

$$\begin{aligned}
\frac{Z_H}{M_{W0}^2/M_W^2} &= 1 \;+\; \frac{1}{8}\frac{g^2}{16\,\pi^2}\frac{m_H^2}{M_W^2}\left(13 - 2\pi\sqrt{3}\right) \\
&+ \frac{1}{16}\left(\frac{g^2}{16\,\pi^2}\frac{m_H^2}{M_W^2}\right)^2\left(3 - 63\,\zeta(3) - \frac{169\,\pi\,\sqrt{3}}{4} + \frac{85\,\pi^2}{12} + 39\,\pi\,C + 57\,\sqrt{3}\,C\right).
\end{aligned} \tag{4.8}$$

Again, we find complete agreement with the numerical result [5] and exact agreement with the result [6], taking into account that their numeric constant K_5 is just minus our integral (3.5).

Acknowledgments

G.J. is grateful to J.J. van der Bij and A. Ghinculov for valuable discussions. This work was supported in part by the Alexander von Humboldt Foundation and the Russian Foundation for Basic Research grant 96-02-19-464.

References

[1] A. Blondel, Plenary talk at the International Conference on High Energy Physics, Warsaw, 1996.

[2] P. Langacker, talk at International Workshop on Supersymmetry and Unification of Fundamental Interactions (SUSY 95), Palaiseau, France, 15-19 May 1995, NSF-ITP-95-140, October 1995, hep-ph/9511207;
W. Hollik, talk at 11th Topical Workshop on Proton-Antiproton Collider Physics (PBARP 96), Padua, Abano Terme, Italy, 26 May - 1 June 1996, Karlsruhe University preprint No KA-TP-19-1996, hep-ph/9608325;
G. Passarino, talk at CRAD96, Cracow, August 1996, hep-ph/9604344.

[3] L. Durand, P.N. Maher and K. Riesselmann, *Phys. Rev.* **D48** (1993) 1084;
K. Riesselmann, *Phys. Rev.* **D53** (1996) 6226;
K. Riesselmann and S. Willenbrock, Reports ILL-(TH)-96-4, TUM-HEP-236/96, August 1996, hep-ph/9608280.

[4] A. Ghinculov and J.J. van der Bij, *Nucl. Phys.* **B436** (1995) 30.

[5] A. Ghinculov, *Phys. Lett.* **B337** (1994) 137; **B346** (1995) 426 (E).

[6] L. Durand, B.A. Kniehl and K. Riesselmann, *Phys. Rev. Lett.* **72** (1994) 2534; **74** (1995) 1699 (E); *Phys. Rev.* **D51** (1995) 5007.

[7] A. Ghinculov, *Nucl. Phys.* **B455** (1995) 21.

[8] A. Frink, B.A. Kniehl, D. Kreimer and K. Riesselmann, *Phys. Rev.* **D54** (1996) 4548.

[9] P.N. Maher, L. Durand and K. Riesselmann, *Phys. Rev.* **D48** (1993) 1061; **D52** (1995) 553 (E).

[10] P. Breitenlohner and D. Mason, *Comm. Math. Phys.* **52** (1977) 11, 39, 55.

[11] R. Scharf and J.B. Tausk, *Nucl. Phys.* **B412** (1994) 523.

[12] D.J. Broadhurst, *Z. Phys.* **47** (1990) 115.

[13] S. Bauberger and M. Böhm, *Nucl. Phys.* **445** (1995) 25.

[14] A.I. Davydychev and J.B. Tausk, *Nucl. Phys.* **B397** (1993) 123.

[15] A. Ghinculov, Freiburg-THEP 96/6, April 1996, hep-ph/9604333.

[16] A.V. Kotikov, *Mod. Phys. Lett.* **A6** (1991) 677, 3133; *Phys. Lett.* **B254** (1991) 158; *Phys. Lett.* **B267** (1991) 123.

[17] F.V. Tkachov, *Phys. Lett.* **B124** (1983) 212; *Int. J. Mod. Phys.* **A8** (1993) 2047.

[18] J. van der Bij, M. Veltman, *Nucl. Phys.* **B231** (1984) 205.